

**POPULATION, SEX-RATIO ESTIMATION OF *LABEO DYOCHAILUS*  
(McClelland) FROM WEST RAM GANGA RIVER, AT CHAUKHUTIA IN  
KUMAUN HILLS**

**RAKESH VERMA & KUMKUM SHAH**

Fisheries Development & Research Laboratory, Department of Zoology  
L.S.M. Govt. P.G. College Pithoragarh, Uttarakhand

**ABSTRACT**

The present paper deals with the sex population of a major carp, *Labeo dyocheilus* (Mc.Cl.) from spring fed river West Ramganga from Kumaun hills, Uttarakhand. The Latitude and Longitude of sampling area is 29° 53' 55" North and 79° 21' 22" East. Total 202 specimens are studied during Sep 2009 to Aug 2011. Sex-ratio was calculated and its significance tested by Chi-square test ( $\chi^2$ ). The population of *Labeo dyocheilus* was found in significant sex ratio 1.77male:1.00 female.

**Key words:** *Labeo dyocheilus*, Sex ratio, Kumaun hills.

**REFERENCES**

- Gaigher, I.G. 1983. Reproduction of *Labeo umbratus* (Pisces, Cyprinidae) in Wuras Dam, a shallow, turbid impoundment. *South African Journal of Zoology* 19: 105-108.
- Holick, J., Hensel, K., Nielslanik, J. and Skace, I.J.1988. The *Eurassian hucken* and *Hucho hucho* largest-salmon of world. Published by Dr.W.Junk. Dordrecht J. Boston: Lancaster, pp... + 239.
- Ikpi,G.U. and Okey. I. B .(2010). Estimation of Dietary Composition and Fecundity of African Carp, *Labeo Coubie*, Cross River, Nigeria. *J. Appl. Sci. Environ. Manage.* Vol. 14 (4) 19 - 24
- Jameela Beevi, K.S. and Ramachandran, A. 2005. Sex ratio in *Puntius vittatus* Day in the fresh water bodies of Ernakulam District, Kerala. *Zoos Print Journal* 20(9): 1989-1990.
- Kumar, K., Bisht,K.L., Dobriyal, A.K., Joshi, H.K., Bahuguna,P.K., Goswami, S., Balodi,V.P. and Thapliyal, A. 2006. Fecundity and sex ratio in a rare hill-stream fish *Botia dayi* Hora from Garhwal Himalaya, Uttranchal. *Uttar Pradesh J. Zool.* 26 (3): 271-276.
- Montchowui, E., Laleye, P., Poncin,P. and Philippart. 2010. Reproductive strategy of *Labeo senegalensis Valenciennes 1842* (Teleostei: Cyprinidae) in the Queme basin, Benin. *African J. Aquatic Science.*Vol.35(1):81-85.
- Nikolsky, G.V.1956. Ecology of fishes. Academic Press, London, pp1-352 (1956).
- Nikolsky, G.V.1980. Theory of fish population dynamics. Bishen singh and Mahendar Pal Singh, India and Ottokoeltz Science Publishers (West Germany), pp317.

- Panwar, B.A. and Mani, U.H. 2006. Sex ratio of *Macrones bleekeri* (Blecker) from Sadatpur lake, Ahmednager, District Maharashtra. *J. Aqua. Biol.* 21(2): 182-185.
- Pathani, S.S. 2000. Reproductive biology of the golden mahseer, *Tor putitora* (Ham.) from Kumaun. *Cold water aquaculture and fisheries*. (Eds H. R. Singh and W.S. Lakra), Narendra Publishing House, Delhi. pp: 253-264.
- Shendge, A.N. and Mani, U.H. 2009: Sex ratio of cyprinidae fish *Cirrhina reba* (Hamilton). *Uttar Pradesh J. Zool.* 29 (2): 217-220.
- Sobhana, B. and Nair, N.B. 1976. Observation on the maturation and spawning of *Puntius sarana subnasutus* (Valenciennes). *Indian J. Fish.* 21(2): 357-359.
- Van Zyl, B.J., Hay, C.J. and Steyn, G.J. 1995. Some aspects of the reproduction biology of *Labeo capensis* (Smith, 1941) (Pisces: Cyprinidae) in relation to exploitation and extreme environmental conditions in Hardap Dam, Namibia. *South African Journal of Aquatic Science* **21**: 88-98.
- Weyl, O.L.F. and Booth, A. J. 1999. On the life history of a cyprinid fish, *Labeo cylindricus*. *Environ. Biol. Fishes* 55:215-225.