

A GENERAL FIXED POINT THEOREM FOR INTEGRAL TYPE IMPLICIT RELATIONS

U.C. GAIROLA AND A.S. RAWAT

Department of Mathematics

H.N.B. Garhwal University, Campus Pauri, Pauri Garhwal-246001, India

Email ucgairola@rediffmail.com and anup_singhrawat@rediffmail.com

Received: 22.09.2012

Revised: 03.11.2012

Accepted: 14.12.2012

ABSTRACT

In this paper we prove a general fixed point theorem for two pairs of mapping satisfying integral type implicit relation on two metric spaces. Our result generalize and extend certain results of Fisher-Murthy [4] and Popa [5].

AMS (MOS) Subject Classifications (2000): 54H25, 47H10.

Key words: Integral type implicit relations, complete metric space.

REFERENCE

- A. Branciari, A fixed point theorem for mappings satisfying a general contractive condition of integral type, *Int. J. Math. Math. Sci.*, 29(9) (2002), 531-536.
- B. Fisher, Fixed point on two metric spaces, *Glasnik Mathem.* 16(36) (1981), 333-337.
- B. Fisher, Related fixed points on two metric spaces, *Math. Sci. Notes, Kobe Univ.*, 10(1982),17-26.
- B. Fisher and P.P. Murthy, Related fixed point theorems for two pairs of mappings on two metric spaces, *Kyungpook Math. J.*, 37 (1997), 343-347.
- V. Popa, A general fixed point theorem for two pairs of mappings on two metric spaces, *Novi. Sad. J. Math.*, 35(2) (2005), 79-83.