



Shear Free Radiating Fluid Ball Without Horizon

Jyoti Rani¹ • Kali Charan^{2*}

¹Department of Mathematics, Uttarakhand Open University, Haldwani, Nainital

²Department of Mathematics, SSJ University, Campus, Almora, 263601.

*Corresponding Author Email id: kcyadav2008@gmail.com

Received: 12.05.2024; Revised: 20.06.2024; Accepted: 21.06.2024

©Society for Himalayan Action Research and Development

Abstract: This paper investigates the novel explicit model of the relativistic field equations for radiating fluid ball in the absence of horizon. The inner metric is without shear, symmetry like sphere and undergoes heat flow in radial direction. The inner metric maintains all appropriate thermodynamic and physical parameters analytically as well as graphically and are matched to the Vaidya external metric across the boundary. Here in the present model, the collapse begins with unlimited mass and radius, and time approaches zero without the event horizon formation. The luminosity is independent of time, and the rate of collapse process will never reach the horizon.

Keywords: radiating fluid ball • exact model • naked singularity • gravitational collapse.

1. Introduction

The gravitational collapse is a theoretical and observable feature in the universe; it captures the focus of general relativity researchers. In relativistic astrophysics the final outcome of the gravitational collapse is an open problem Joshi and Malafarina (2011). As of right now, no established theory exists that can predict this collapse. However, there are many contrary situations where the creation of a naked singularity is most likely to occur.

Constructing a realistically model of a collapsing structure is essential in the context of fundamental gravity in order to fully comprehend the system and the physical behavior of the decreasing structure. Because the laws underlying field equations are highly nonlinear, this is an extremely challenging task to solve them. A number of attempts have been made to make easier things, and different approaches are regularly used to do so. The important contribution is made by Oppenheimer and Snyder (1939), which examined the collapse

of an incredibly idealized spherically statuesque particles cloud, was the first step in this direction.

In an effort to comprehend the features and nature of collapsing things, numerous strive have been formed since then to build accurate models of collapsing system. It acquired enormous momentum when Vaidya (1951) offered a solution that combined the modified equations for an adiabatic dispensation of matter introduced by Misner (1965) and Lindquist et al. (1965) with a model defining the external gravitational expansion of a star objects with radiation leaving it.

Collapsing of an object is recognized as a very dissipative form of energy (Herrera and Santos 2004, Herrera et al. 2006, Mitra 2006). On the other hand, two limiting scenarios are outlined where energy dissipation via collapsing fluid distribution occurs. In the first situation, Tewari (1988), Pant et al. (1990), Tewari (1994) discusses a number of radiating matter solutions. Additionally, Pant and Tewari (2011) offered a



quasar structure that was suitable to every extent. As opposed to the second, which is the diffusion case, where dissipation is represented by a heat flow. In this direction Glass (1981) and Santos (1985) with reference to the junction conditions of a radially heated collapsing system are significant. The numerous papers on related field have been published by Banerjee et al. (2002), Maharaj et al. (2005), de Oliveira et al. (1985), Maiti (1987), de Oliveira-Santos (1987), Sarwe-Tikekar (2010), Bonnor et al. (1989), Tewari (2012,2013), Pinheiro-Chan (2013), Tewari et al. (2014, 2015a, 2015b, 2015c, 2015d), Ivanov (2012), Charan et al. (2021), Kauntye Acharya et al. (2023), Ghosh Avisikta (2024) and references within that describe the energy-radiating collapse of a collapsing fluid. Govender et al. (2020) present realistic analytic models characterized by only two metric

components, constrained by spatial development limited through isotropy conditions and the radiating ideal fluid with heat flow is typically explored in order to explain the calculations. In considering the aforementioned considerations, we examine Tewari et al. (2014) solution in an effort to create a realistic radiating ball model. Over the boundary, the Vaidya exterior line element (1953) is matched with the interior line element.

This study is organized into five sections. In section 2, junction circumstances and space-time are discussed. In Section 3 using method of variable separation to separate the space and time variables. The model's accompanying energy conditions are covered in Section 4. In Section 5, the graphical representation and evaluation of the energy conditions are discussed.

2. The matching conditions

The metric element with time like space is given by the following relation

$$ds^2 = -X^2(r,t)dt^2 + Y^2(r,t)\{dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2)\} \quad (1)$$

For a fluid, the energy-momentum tensor is given as

$$T_{ij} = (\epsilon + p)u_i u_j + p g_{ij} + q_i u_j + q_j u_i \quad (2)$$

where symbols have their usual meanings.

For fluid distribution (1), the fluid collapse rate is determined by

$$\theta = \frac{3\dot{Y}}{XY} \quad (3)$$

Given (1) and (2), the system of equations that follows

$$\kappa\epsilon = -\frac{1}{Y^2}\left(\frac{2Y''}{Y} - \frac{Y'^2}{Y^2} + \frac{4Y'}{rY}\right) + \frac{3\dot{Y}^2}{X^2Y^2} \quad (4)$$

$$\begin{aligned} \kappa p = & -\frac{1}{Y^2}\left(\frac{2Y'^2}{Y^2} + \frac{2XY'}{XY} + \frac{2X'}{XY} + \frac{2X'}{rX} + \frac{2Y'}{rY}\right) \\ & + \frac{1}{X^2}\left(-\frac{2\ddot{Y}}{Y} - \frac{\dot{Y}}{Y^2} + \frac{2\dot{X}\dot{Y}}{XY}\right) \end{aligned} \quad (5)$$

$$\kappa p = -\frac{1}{Y^2}\left(\frac{2Y''}{Y} - \frac{Y'^2}{Y^2} + \frac{X'}{rX} + \frac{X''}{X} + \frac{Y'}{rY}\right) + \frac{1}{X^2}\left(-\frac{2\ddot{Y}}{Y} - \frac{\dot{Y}}{Y^2} + \frac{2\dot{X}\dot{Y}}{XY}\right) \quad (6)$$

$$\kappa q = \frac{-2}{XY^2}\left(-\frac{\dot{Y}}{Y} + \frac{Y'\dot{Y}}{Y^2} + \frac{X'\dot{Y}}{XY}\right) \quad (7)$$

The external metric is given by (1953) as

$$ds^2_{\pm} = -\left(1 - \frac{2M(v)}{R}\right)dv^2 - 2dRdv + R^2(d\theta^2 + \sin^2\theta d\phi^2) \quad (8)$$



The matching of (1) and (8) at the boundary is given by Santos (1985) as

$$(rY)_{\Sigma} = R_{\Sigma}v = R(r) \quad (9)$$

$$(p_r)_{\Sigma} = (qY)_{\Sigma} \quad (10)$$

$$m_{\Sigma}(r, t) = M(v) = \left(\frac{r^3 Y \dot{Y}^2}{2X^2} - r^2 Y' - \frac{r^3 Y^2}{2Y} \right)_{\Sigma} \quad (11)$$

This mass at the boundary is given by Cahill et al. (1970) and Misner et al. (1964).

The Z_{Σ} and L_{Σ} on the boundary are obtained from the following relations

$$L_{\Sigma} = \frac{1}{2} \{ r^2 Y^3 q \}_{\Sigma} \quad (12)$$

$$z_{\Sigma} = \left[1 + \frac{rY'}{Y} + \frac{r\dot{Y}}{X} \right]_{\Sigma}^{-1} - 1 \quad (13)$$

The L_{∞} at rest is given as

$$L_{\infty} = - \frac{dM}{dv} = \frac{L_{\Sigma}}{(1+z_{\Sigma})^2} \quad (14)$$

3. Solution of field equations

In equation (1), taking into consideration the following different forms for the metric

$$X(r, t) = X_0(r)g(t) \quad (15)$$

$$Y(r, t) = Y_0(r)f(t) \quad (16)$$

With the help of field equations (4), (5), (7), (15) and (16) we have

$$\epsilon = - \frac{1}{f^2 Y_0'} \left(\frac{2Y_0'^2}{Y_0} - \frac{Y_0^2}{Y_0'} + \frac{4Y_0'}{rY_0} \right) + \frac{3\dot{f}^2}{X_0^2 g^2 f^2} \quad (17)$$

$$p = \frac{1}{Y_0'} \left(\frac{Y_0'^2}{Y_0} + \frac{2Y_0'}{rY_0} + \frac{2X_0'Y_0'}{X_0Y_0} + \frac{2X_0'}{rX_0} \right) + \frac{1}{X_0^2 g^2} \left(- \frac{2\dot{f}}{f} - \frac{\dot{f}^2}{f^2} \right) \quad (18)$$

$$q = - \frac{2X_0'\dot{f}}{X_0^2 Y_0' f^3} \quad (19)$$

The boundary conditions (10), $(p_r)_{\Sigma} = (qY)_{\Sigma}$ now yields at $r = r_{\Sigma} = R_{\Sigma}$ in view of (16), (18) and (19)

$$\ddot{f} + \dot{f}^2 - 2\eta\dot{f} = \zeta \quad (20)$$

where η and ζ are constant and given by

$$\eta = \left(\frac{X_0'}{Y_0} \right) \quad (21)$$

$$\zeta = \frac{X_0^2}{Y_0^2} \left(\frac{Y_0'^2}{Y_0} + \frac{2Y_0'}{rY_0} + \frac{X_0'Y_0'}{X_0Y_0} + \frac{2X_0'}{rX_0} \right)_{\Sigma} \quad (22)$$

In view of particular solution of above equation is given by

$$f = - \alpha t \quad (23)$$

where α is arbitrary constant.

In order for the configuration to collapse we must have $\alpha < 0$, meaning that α must be positive.

$$\alpha = - \eta + \sqrt{\eta^2 + \zeta}, \quad \zeta > 0 \quad (24)$$

The isotropy pressure would given as below

$$\frac{X_0''}{X_0} + \frac{Y_0''}{Y_0} = \left(\frac{2Y_0'}{Y_0} + \frac{1}{r} \right) \left(\frac{X_0'}{X_0} + \frac{Y_0'}{Y_0} \right) \quad (25)$$



The solution of (25) given by Tewari et al. (2014)

$$X_0 = d_2(1 + c_1 r^2)^{\frac{n}{l+1}} + d_1(1 + c_1 r^2)^{\frac{2-n}{l+1}+1} \quad (26)$$

$$Y_0 = c_2(1 + c_1 r^2)^{\frac{1}{l+1}} \quad (27)$$

$$n = \frac{1}{2} \left\{ (l+3) \pm (l^2 + 10l + 17)^{\frac{1}{2}} \right\} \quad (28)$$

Tewari's variety of solutions (2014) is given by equations (26) and (27) for various values of n and l . To create a new realistic model, we take the positive constant $c_1 = 1, d_1 + d_2 = 1$, and assume $l = -5 + 2\sqrt{2}$ in the context of (26) and (27) and obtain

$$X_0 = (1 + c_1 r^2)^{-\frac{\sqrt{2}}{4}} \quad (29)$$

$$Y_0 = (1 + c_1 r^2)^{-\frac{2-\sqrt{2}}{4}} \quad (30)$$

The expression for pressure, density and heat flux can be obtained as

$$\epsilon = \frac{1}{t^2(1+c_1 r^2)^{-\frac{\sqrt{2}}{2}}} \left[\frac{c_1 \{6\sqrt{2}(\sqrt{2}+1) + c_1 r^2\}}{2\alpha^2(1+c_1 r^2)} - 1 \right] \quad (31)$$

$$p = \frac{1}{t^2(1+c_1 r^2)^{-\frac{\sqrt{2}}{2}}} \left[\frac{c_1 \{-4(\sqrt{2}+1) + c_1 r^2\}}{2\alpha^2(1+c_1 r^2)} - 1 \right] \quad (32)$$

$$q = \frac{\sqrt{2}c_1 r}{\alpha^2 t^3(1+c_1 r^2)^{-\frac{3\sqrt{2}}{4}}} \quad (33)$$

From the above equations, we obtain

$$\eta = -\frac{\sqrt{2}c_1 r}{2(1+c_1 r^2)^{\frac{1}{2}}} \quad (34)$$

$$\zeta = \frac{c_1 \{-4(\sqrt{2}+1) + c_1 r^2\}}{2(1+c_1 r^2)} \quad (35)$$

In the solution $f = -\alpha t$, the constant α chosen to be positive ($\eta > 0, \zeta > 0$) so the collapsing phase corresponds to $-\infty < t < 0$. The isotropic pressure p , density ϵ and q have value 0 at $t \rightarrow \infty$ and diverge when $t \rightarrow 0$. The matter density and isotropic pressure evolve with time as t^{-2} while q evolves as t^{-3} .

4. Energy conditions

The $\epsilon > 0, p > 0$ are satisfied if

$$\alpha^2 < \frac{c_1 \{-4(\sqrt{2}+1) + c_1 r^2\}}{2(1+c_1 r^2)} \quad (36)$$

and $\epsilon' < 0, p' < 0$ are satisfied if

$$\alpha^2 < \frac{c_1 \{(4+\sqrt{2}) + c_1 r^2\}}{2(1+c_1 r^2)} \quad (37)$$

Above conditions (36) and (37) are satisfied if $\alpha^2 < c_1 \{-2(\sqrt{2} + 1)\}$.

Now $(\epsilon - p) > 0$ are satisfied if

$$\alpha^2 < \frac{-c_1 \{(5\sqrt{2}+8)\}}{4(1+c_1 r^2)} \quad (38)$$



$(\epsilon - 3p) > 0$ are satisfied if

$$\alpha^2 < \frac{c_1\{-3(4+3\sqrt{2})+c_1r^2\}}{6(1+c_1r^2)} \quad (39)$$

Then all the physical suitable conditions will be satisfied if $\frac{c_1\{-3(4+3\sqrt{2})+c_1r^2\}}{6(1+c_1r^2)} < \alpha^2 < c_1\{-2(\sqrt{2}+1)\}$. Due to radial heat flow, the fluid must satisfy the condition

$(\epsilon + p) > 2|q|$, where $|q| = (g_{ij}q^i q^j)^{\frac{1}{2}}$, to ensure consistency with all energy conditions. This would require,

$$\left[\frac{c_1\{(4+\sqrt{2})+c_1r^2\}}{2(1+c_1r^2)} + 1 \right] > -\frac{\sqrt{2}c_1r}{\alpha(1+c_1r^2)^{\frac{1}{2}}} \quad (40)$$

which would always be true as it could be written as

$$\left[\sqrt{2} + \frac{c_1r}{\alpha(1+c_1r^2)^{\frac{1}{2}}} \right]^2 > \frac{-c_1(4+\sqrt{2})}{\alpha^2(1+c_1r^2)} \quad (41)$$

Now we obtained the explicit formulation for surface luminosity radius, surface luminosity, total energy entrapped inside Σ as

$$R_{\Sigma}(v) = (rY)_{\Sigma} = -r_{\Sigma}\alpha t (1 + c_1r_{\Sigma}^2)^{\frac{-2-\sqrt{2}}{4}} \quad (42)$$

$$M(v) = -\frac{1}{2}atr_{\Sigma}^3(1 + c_1r_{\Sigma}^2)^{\frac{-6-\sqrt{2}}{4}} \left[\alpha^2 + \frac{c_1\{(4-2\sqrt{2})-c_1r^2\}}{2(1+c_1r^2)} \right] \quad (43)$$

In view of (3), (12), (13) and (14), we get

$$\Theta = \frac{3}{t(1+c_1r_{\Sigma}^2)^{\frac{-\sqrt{2}}{4}}} \quad (44)$$

$$L_{\Sigma} = \frac{-\sqrt{2}\alpha c_1 r_{\Sigma}^2}{2(1+c_1r_{\Sigma}^2)^{\frac{3}{2}}} \quad (45)$$

$$L_{\infty} = \frac{-\sqrt{2}\alpha c_1 r_{\Sigma}^2}{2(1+c_1r_{\Sigma}^2)^{\frac{3}{2}}} \left[1 - \frac{\sqrt{2}}{2} c_1 r_{\Sigma}^2 - r_{\Sigma} \alpha (1 + c_1 r_{\Sigma}^2)^{\frac{1}{2}} \right]^2 \quad (46)$$

$$z_{\Sigma} = \frac{(1+c_1r_{\Sigma}^2)}{\left[1 - \frac{\sqrt{2}}{2} c_1 r_{\Sigma}^2 - r_{\Sigma} \alpha (1 + c_1 r_{\Sigma}^2)^{\frac{1}{2}} \right]} - 1 \quad (47)$$

To avoid the situation of horizon appearance the model parameters might be set that $\frac{2M}{R_{\Sigma}} < 1$. This implies

that $\alpha^2 < \frac{\left(1 - \frac{\sqrt{2}}{2} c_1 r_{\Sigma}^2\right)^2}{r_{\Sigma}^2(1+c_1r_{\Sigma}^2)}$.

The analysis of temperature regarding the model is obtained from the following law given by Maartens (1995), Israel et al. (1979) and Martinez (1996).

$$\tau(g^{ij} + u^i u^j) u^{\alpha} q_{j\alpha} + q^j = -\mathbb{K}(g^{ij} + u^i u^j) [T_j + T u_j] \quad (48)$$

where symbols have their usual meanings.

The relaxation time in above expression (48) is given by the following relation

$$q = -\mathbb{K} \frac{1}{r^2 f^2} \left(T' + T \frac{\dot{X}_0}{X_0} \right) = -\frac{2X_0 \dot{f}}{X_0^2 r_0^2 f^2} \quad (49)$$



The effective surface temperature given as Schwarzschild (1958) and could be evaluated with the help of following relation

$$T_{\Sigma}^4 = \left\{ \frac{1}{\pi \delta (r Y_0 f)^2} \right\}_{\Sigma} L_{\infty} = \frac{\sqrt{2} c_1 r_{\Sigma}}{2 \pi \delta a t^2 (1 + c_1 r^2)^{\frac{5 + \sqrt{2}}{2}}} \left[1 - \frac{\sqrt{2}}{2} c_1 r_{\Sigma}^2 - r_{\Sigma} \alpha (1 + c_1 r_{\Sigma}^2)^{\frac{1}{2}} \right]^2 \quad (50)$$

where $\delta = \frac{\pi^2 k^4}{15 \hbar^3}$.

Choosing $\Omega = 3$ in Misner et al. (1964), the function T_0 is given by

$$T_0(t) = \frac{1}{a t^2} \left[\frac{8}{3 \kappa \gamma (1 + c_1 r^2)^{\frac{3\sqrt{2}}{4}}} + \frac{\sqrt{2} c_1}{\pi \delta r^2 a t (1 + c_1 r^2)^{\frac{1 + \sqrt{2}}{2}}} \frac{1}{(1 + \Sigma)^2} \right]_{\Sigma} \quad (51)$$

The temperature inside the star is given by

$$T^4 = \left[\frac{T_0(t)}{(1 + c_1 r^2)^{-\sqrt{2}}} \right] - \left[\frac{8}{3 \gamma t (1 + c_1 r^2)^{\frac{\sqrt{2}}{4}}} \right] \quad (52)$$

5. Graphical representations of model parameters

We have taken model parameters as following for pictorial representation

$c_1 = 0.1, r_{\Sigma} = 0.5$.




Fig.1 Evolution of Energy density versus radial and temporal coordinates




Fig. 2 Evolution of pressure versus radial and temporal coordinates




Fig. 3 Evolution of Heat flux versus radial and temporal coordinates




Fig. 4 Evolution of Collapse rate versus radial and temporal coordinates




Fig. 5 Evolution of Mass energy versus temporal coordinates




Fig.6 Evolution of luminosity radius versus temporal coordinates




Fig.7 Evolution of Surface temperature versus temporal coordinates

6. Conclusion

The exact model of shear-free spherically symmetric isotropic fluid distributions of matter without the appearance of horizon are presented in this study. By assuming $n = -1 + \sqrt{2}$, we create a new horizon-free model. When time

approaches zero, the surface temperature is zero, yet it progressively rises and eventually diverges. The luminance for a distant observer has been shown as time independent in this model. The model fulfilled all the basic properties of a collapsing radiating star. The rate



of collapse grows from the beginning condition in a realistic model proposed by Tewari et al. (2015a) and the mass-to-radius ratio is time independent. Finally, the rate of mass loss was balanced by the collapse of boundary radius; as a result, our model did not form in the horizon, and it was observed that the boundary surface could never reach the horizon when

$$\alpha^2 < \left(1 - \frac{\sqrt{2}}{2} c_1 r_{\frac{\Sigma}{2}}^2\right)^2 / r_{\frac{\Sigma}{2}}^2 (1 + c_1 r_{\frac{\Sigma}{2}}^2).$$

References

- Acharya Kauntey, Pandey Kshitij, Bambhaniya Parth, Joshi, Pankaj S and Patel Vishva (2023). Naked Singularity as a Possible Source of Ultra-High Energy Cosmic Rays, arXiv:2303.16590v1[gr-qc].
- Banerjee A, Chatterjee S and Dadhich N (2002). Spherical Collapse with Heat Flow and without and without Horizon, *Modern Physics Letters A*, 17: 2335-2339.
- Bonnor WB, De Oliveira A, and Santos NO (1989). Radiating spherical collapse, *Phys. Rep.*, 181(5):269-326.
- Cahill ME and McVittie GC (1970). Spherical Symmetry and Mass- Energy in General Relativity, *I. General Theory*, *J. Math. Phys.*, 11:1382-1391.
- Charan Kali, Yadav Om Prakash and Tewari, BC (2021). Charged anisotropic spherical collapse with heat flow, *Eur.Phys.J.C.*, 81:60.
- de Oliviera, AKG and Santos NO (1987). Non-adiabatic Gravitational Collapse, *Astrophys.J.* 312: 640-645.
- de Oliviera, AKG, Santos NO and Kolassis CA (1985). Collapse of Radiating Star, *Monthly Notices of the Royal Astronomical Society*, 216: 1001-1011.
- Ghosh Avisikta (2024). Study of Orbital Dynamics in Singular and Regular Naked Singularity Space-times, *arXiv* 2401: 1077[gr-qc].
- Glass, EN (1981). Shear-free collapse with heat flow, *Phys. Lett. A*, 86: 351-352.
- Govendar M, Maharaj A, Singh, Ksh Newton and Neeraj Pant (2020). Dissipative Collapse of a Karmarkar Star, *Modern Physics Letters A.*, 35(20): 2050164.
- Herrera L and Santos NO (2004). Dynamics of dissipative gravitational collapse, *Phys.Rev.D*, 70:084004,1-8.
- Herrera L, Di Prisco A, Martin J and Ospino J (2006). Some analytical models of radiating collapsing spheres, *Phys,Rev D.*, 74: 044001 (2006).
- Israel W and Stewart J (1979). Transient relativistic thermodynamics and kinetic theory, *Annals of Physics*, 118: 341-372.
- Ivaniov BV (2012). Collapsing shear-free perfect fluid spheres with heat flow, *General Relativity and Gravitation* 44: 1835-1855.
- Joshi PS and Malafarina D (2011). Recent developments in Gravitational Collapse and Space-time singularities, *International Journal of modern physics D*, 20: 2641-2729.
- Lindquist JM, Plotkin SA, Shaw L, Silden RV and Williams ML (1965). Congenital rubella syndrome as a systemic infection. Studies of affected infants born in Philadelphia, *U.S.A Br Med J*, 25475.
- Maartens R (1995). Dissipative cosmology, *Classical and Quantum Gravity*, 12: 1455-1465.
- Mahraj SD and Govender M (2005). Radiating Collapse with Vanishing Weyl Stresses, *International Journal of Modern Physics D*, 14: 667-676.
- Maiti SR (1987). Fluid with heat flux in a conformally flat space-time, *Phys.Rev.D.*, 25: 2518.



- Martinez J (1996). Transport processes in the gravitational collapse of an anisotropic fluid, *Physical Review D*, 53: 6921-6940.
- Misner CW (1965). Relativistic equations for spherical gravitational collapse with escaping Neutrinos. *Phys. Rev.*, 137(5B): B1360.
- Misner CW and Sharp David H (1964). Relativistic Equations for Adiabatic, Spherically Symmetric Gravitational Collapse, *Phys. Rev.B.*, 136: B571-B576.
- Mitra, A(2006). Why Gravitational Contraction Must Accompanied by Emission of Radiation Both in Newtonian and Einstein Gravity, *Physical Review D.*, 74: 024010.
- Oppenheimer JR and Snyder H (1939). On Continued gravitational contraction, *Physical Review*, 56:455-459.
- Pant DN and Tewari, BC (1990). Conformally-Flat Matrix Representing a Radiating Fluid Ball, *Astrophys Space Sci.*, 163: 223-227.
- Pant N and Tewari BC (2011). Horizon-free gravitational collapse of radiating fluid sphere, *Astrophys. Space Sci.*, 331(2): 645–650.
- Pinheiro G and Chan R (2013). Radiating Shear-Free Gravitational Collapse with Charge, *General Relativity and Gravitation*, 45: 243-261.
- Santos NO (1985). Non-adiabatic radiating collapse, *MNRAS*, 216: 403-410.
- Sarwe R and Tikekar R (2010). Non-Adiabatic Gravitational Collapse of a Superdense star, *Int.J. Mod.Phys.*, D (19): 1889-1904.
- Schwarzschild, M (1958). Structure and Evaluation of stars, *Dover. New. York*.
- Tewari BC and Charan K (2014). Radiating star, shear-free gravitational collapse without horizon, *Astrophysics. Space Sci.*, 351: 613-617.
- Tewari BC and Charan K (2015b). Gravitational Collapse of Radiating Star with Radial Heat Flow, *International journal of applied mathematical Sci*, 8 (21): 77-80.
- Tewari BC and Charan K(2015a). Dissipative Spherical Gravitational Collapse of Isotropic Fluid, *J Modern Physics*, 6: 453-462.
- Tewari, BC (2012). Relativistic collapsing radiating stars, *Astrophys. Space Sci.* 342:73-77.
- Tewari, BC (2013). Collapsing shear-free radiating fluid spheres, *Gen. Relativ. Gravit.*, 45: 1547-1558.
- Tewari, BC and Charan K (2015c). Horizon free eternally collapsing anisotropic radiating star, *Astrophys Space Sci.* 357:107.
- Tewari, BC and Charan K (2015d). Gravitational Collapse, shear free anisotropic radiating star, Gravitational Collapse, shear free anisotropic radiating star, *General Relativity and Quantum Cosmology*, arXiv: 1503.02165 [gr-qc].
- Tewari, BC (1988). Radiating Fluid spheres in General Relativity, *Astrophysics and Space Science*, 149: 233-239.
- Tewari, BC (1994). Relativistic Radiating Fluid Distribution, *Indian Journal of Pure and Applied Physics*, 32: 504-507.
- Vaidya PC (1951). The gravitational field of a radiating star, *Proc. Indian Acad. Sci., Sect. A, Phys. Sci.*, 33:264-276.
- Vaidya PC (1953). Newtonian time in general relativity, *Nature*, 171: 60-261.